



A Primary Intraosseous Rhabdomyosarcoma of Mandible: Report of a Rare Case and Review of Literature

Shabnam Mashhadi^{1*}, Ata Garajei², Pouyan Aminishakib^{1,3}

1. Department of Pathology, Cancer Institute, Imam Khomeini Hospital Complex, Tehran University of Medical Sciences, Tehran, Iran
2. Department of Head and Neck Surgical Oncology and Reconstructive Surgery, Cancer Institute, Imam Khomeini Hospital Complex, Tehran University of Medical Sciences, Tehran, Iran
3. Oral and Maxillofacial Pathology Department, School of Dentistry, Tehran University of Medical Sciences, Tehran, Iran

Article Info

Article type:
Case Report

Article History:

Received: 25 Dec 2023

Accepted: 30 May 2024

Published: 18 Jan 2025

* Corresponding author:

Department of Pathology, Cancer Institute, Imam Khomeini Hospital Complex, Tehran University of Medical Sciences, Tehran, Iran

Email: Shabnam.mashadi@gmail.com

ABSTRACT

Rhabdomyosarcoma is a rare soft tissue malignant neoplasm with striated muscle differentiation often found in the head and neck region. In our case a 33-year-old man presented with a bulging mass near his right mandibular second molar tooth, identified as a unilocular destructive lesion on an X-ray. An incisional biopsy confirmed rhabdomyosarcoma, leading to a hemi-mandibulectomy. Histopathologic findings showed spindle and rhabdoid cells with positive immunoreactivity to Desmin, myogenin and myoD1. In conclusion, it is crucial to differentiate rhabdomyosarcoma from other more common intraosseous malignancies, and using Desmin staining as part of the initial immunohistochemical panel can aid in the diagnosis process.

Keywords: Rhabdomyosarcoma; Mandible; Neoplasm

- **Cite this article as:** Mashhadi Sh, Garajei A, Aminishakib P. A Primary Intraosseous Rhabdomyosarcoma of Mandible: Report of a Rare Case and Review of Literature. *Front Dent.* 2025;22:04. <https://doi.org/10.18502/fid.v22i4.17543>

INTRODUCTION

Rhabdomyosarcoma is a rare malignant soft tissue neoplasm that originates from mesenchymal cells with striated muscle cell differentiation [1]. It accounts for only 1% of all malignancies, with 35% of cases occurring in the head and neck region [2,3]. Oral cavity rhabdomyosarcoma is even rarer, representing 10-12% of cases of this type of sarcoma [4]. It is commonly found in the tongue, soft palate, and hard palate and typically affects younger individuals [2,4]. Here, we present a case of an adult patient with a mandibular mass diagnosed

as primary intraosseous rhabdomyosarcoma.

CASE REPORT

A 33-year-old man presented with a swollen and ulcerated area in his oral cavity in October 2020. His right mandibular second molar tooth was extracted, and he remained symptom-free until January 2021 when a bulge developed in his right mandibular region. An incisional biopsy revealed a spindle cell neoplasm with positive immunoreactivity for Vimentin and Desmin stains and negative reactions for CD31, CD34, CD56, CD99 and

S100. The patient was referred to the Cancer Institute Hospital, Tehran, Iran for further evaluation. Imaging studies showed a destructive unilocular radiolucent lesion extending from the alveolar ridge to the inferior border of the mandibular body involving both buccal and lingual plates with perforation and soft tissue involvement. The lesion was in close proximity to the inferior alveolar nerve (Fig 1).



Fig 1. Panoramic view of the patient. The destructive lesion is seen at right mandibular region.

An excisional surgery was performed, and the specimen revealed a well-circumscribed mass measuring 4x2.5x2.5cm on the superior aspect of the mandible composed of small round rhabdoid cells and spindle cells with hyperchromatic nuclei and prominent nucleoli with few mitotic figures (Fig 2). Immunohistochemical staining confirmed the primary diagnosis of intraosseous rhabdomyosarcoma, spindle cell variant. The neoplastic cells were negative for S100, SOX10, and CD34 but positive for Desmin, Myogenin, and Myo-D1 (Fig 3). The proliferation marker Ki-67 showed a proliferative activity of 20% in neoplastic cells.

DISCUSSION

Rhabdomyosarcoma is rarely seen in adults, and its etiology remains unclear [5]. It presents clinically with a wide range of symptoms, from a cutaneous facial nodule to a progressed painless facial swelling [6]. Histologically, four subtypes of this soft tissue tumor are described: alveolar, embryonal (the most common), botryoid and spindle cell. The spindle cell variant is the least common but has a better prognosis [7]. It is believed to originate from primitive mesenchymal cells, despite showing skeletal muscle differentiation [8]. Rhabdomyosarcoma is unlikely to originate from skeletal muscle fibers, as they are long-lasting cells that do not divide after birth. The interosseous origin of this neoplasm in adults has been rarely reported (Table 1). Our case is the fifth reported case of an adult with primary interosseous rhabdomyosarcoma and the third reported case involving the mandible. The differential diagnosis for our case includes locally aggressive odontogenic tumors, odontogenic carcinomas, locally aggressive non-odontogenic tumors, primary interosseous non-odontogenic carcinomas and malignant bone and cartilage forming tumors. Microscopic examination ruled out malignancies with neural differentiation, vascular-origin neoplasms, and sarcomas. Thus, SOX10 and S100 staining were negative, and CD34 staining ruled out vascular differentiation. Positive staining for Desmin, Myogenin, and MyoD1 in the neoplastic cells confirmed the diagnosis of rhabdomyosarcoma.

Table 1. Literature review

Author	Year	Patient's age	Symptoms	location	Histology subtype
Kimario, O.M [9]	2021	25	Facial swelling	Right maxilla	Alveolar type
Rodrigues [10]	2021	33	Asymptomatic gingival mass	Maxilla	Epithelioid
Joy [11]	2018	52	Facial swelling	Left mandible	Spindle cell
Sadhasivam [12]	2018	25	Facial swelling	Mandible	Spindle cell




Fig 2. Hematoxylin and eosin staining reveals a hypercellular round to spindle cell neoplastic lesion with cellular atypia and scattered mitotic figures. a. x40, b. x100, c. x400.




Fig 3. Immunohistochemical staining shows the expression of skeletal muscle antigen in neoplastic cells. a. Desmin X100, b. MyoD1 X100, c. Myogenin X100.

CONCLUSION

Primary intraosseous rhabdomyosarcoma is an extremely rare neoplasm that needs to be distinguished from a long-list of differential diagnoses. It may be advisable to use a Desmin stain as part of the initial immunohistochemical panel as a screening test to confirm or rule out rhabdomyosarcoma.

CONFLICT OF INTEREST STATEMENT

None declared.

REFERENCES

1. Muala HA, Alaboudy A. Embryonic Rhabdomyosarcoma Case Report and Literature Review. *Internet J Head Neck Surg.* 2009;4(2):1-5.
2. Shrutha SP, Vinit GB. Rhabdomyosarcoma in a pediatric patient: A rare case report. *Contemp Clin Dent.* 2015 Jan-Mar;6(1):113-5.
3. Hartmann S, Lessner G, Mentzel T, Kübler AC, Müller-Richter UD. An adult spindle cell rhabdomyosarcoma in the head and neck region with long-term survival: a case report. *J Med Case Rep.* 2014 Jun 19;8:208.
4. Arul AS, Verma S, Arul AS, Verma R. Oral rhabdomyosarcoma-embryonal subtype in an adult: A rarity. *J Nat Sci Biol Med.* 2014 Jan;5(1):222-5.
5. Hettmer S, Li Z, Billin AN, Barr FG, Cornelison DD, Ehrlich AR, et al. Rhabdomyosarcoma: current challenges and their implications for developing therapies. *Cold Spring Harb Perspect Med.* 2014 Nov 3;4(11):a025650.
6. Loduca SV, Mantesso A, de Oliveira EM, de Araújo VC. Intraosseous rhabdomyosarcoma of the mandible: a case report. *Int J Surg Pathol.* 2003 Jan;11(1):57-60.
7. Rudzinski ER, Anderson JR, Hawkins DS, Skapek SX, Parham DM, Teot LA. The World Health Organization Classification of Skeletal Muscle Tumors in Pediatric Rhabdomyosarcoma: A Report From the Children's Oncology Group. *Arch Pathol Lab Med.* 2015 Oct;139(10):1281-7.
8. Sudhakar S, Geethika V, Rao N, Smitha B, Kiran ChS. Alveolar rhabdomyosarcoma on the left maxillary alveolus: a unique presentation. *J Clin Diagn Res.* 2015 Feb;9(2):ZD07-9.
9. Kimario OM, Washington L, Aloyce A,

Mbunda F, Donasiano A, Saitabau Z, et al. Rhabdomyosarcoma in a Female Adult a Rare Case Report at BMC Mwanza. East African Scholars J Med Surg. 2021 Aug;3(8):136-8.

10. Rodrigues-Fernandes CI, Sánchez-Romero C, de Almeida OP, Alves FA, Carvalho EJA, de Castro JFL, et al. Oral epithelioid rhabdomyosarcoma: Report of a rare case and literature review of a distinct variant of rhabdomyosarcoma. Oral Oncol. 2022 Jan;124:105498.

11. Joy T, Tupkari JV, Hanchate AV, Siwach P. Oral rhabdomyosarcoma in an adult male: A rare case report. J Oral Maxillofac Pathol. 2018 May-Aug;22(2):285.

12. Sadhasivam V, Sherwood IA, Gutmann JL, Gururaj N, Mahalakshmi V, Doss DM. Spindle cell rhabdomyosarcoma of the anterior mandible manifesting as a periapical lesion of pulpal origin: Case report. Aust Endod J. 2019 Apr;45(1):129-134.