



# Antimicrobial Activity of the Essential Oil Obtained from the Seed and Oleo-Gum-Resin of *Ferula Assa-Foetida* against Oral Pathogens

Alireza Daneshkazemi<sup>1</sup>, Hengameh Zandi<sup>2</sup>, Abdolrahim Davari<sup>1</sup>, Mahmood Vakili<sup>3</sup>, Majid Emtiazi<sup>4</sup>, Reza Lotfi<sup>5</sup>, Seyed Mohammad Reza Masoumi<sup>5,6\*</sup>

1. Department of Operative Dentistry, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
2. Department of Microbiology, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
3. Department of Community Medicine, Health Monitoring Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
4. Department of Traditional Medicine, College of Persian Medicine, Shahid Sadoughi University of Medical Sciences, Ardakan, Yazd, Iran
5. School of Dentistry, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
6. Student Research Committee, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

## Article Info

**Article type:**  
Original Article

## Article History:

Received: 15 April 2018  
Accepted: 14 October 2018  
Published: 30 April 2019

## \* Corresponding author:

Student Research Committee, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

E-mail: smrmasoumi1994@gmail.com

## ABSTRACT

**Objectives:** The objective of this study was to evaluate the antimicrobial activity of the essential oil obtained from oleo-gum-resin and seeds of *Ferula assa-foetida*.

**Materials and Methods:** *Ferula assa-foetida* plants were collected from Tabas, Yazd Province, Iran, during summer 2017. Then, essential oils were obtained from its seeds and oleo-gum-resin using hydrodistillation. Gas chromatography-mass spectrometry (GC-MS) test was performed to determine the contents of the essential oils. Four different concentrations of each oil were prepared (2.5, 5, 10, and 20 µg/ml), and the antimicrobial activity of each dose against four oral bacteria (*Streptococcus mutans*, *Streptococcus sobrinus*, *Streptococcus sanguis*, *Streptococcus salivarius*, and *Lactobacillus rhamnosus*) was evaluated using the disk diffusion method. The data were analyzed using analysis of variance (ANOVA) and Kruskal-Wallis test in SPSS 17 software.

**Results:** The GC-MS findings exhibited that the main compounds found in essential oils yielded from the seeds and oleo gum resin were (Z) -1-propenyl sec-butyl disulfide and (E) -1-propenyl sec-butyl disulfide. *Ferula assa-foetida* plant showed a significant antimicrobial effect ( $P < 0.05$ ). The essential oil from *Ferula assa-foetida* oleo-gum-resin had significantly stronger antibacterial properties compared to the essential oil from *Ferula assa-foetida* seeds ( $P < 0.001$ ). Both essential oils showed antibacterial properties similar to that of Chlorhexidine. The growth inhibition zone was significantly dependent on the essential oil concentration for all bacteria ( $P < 0.05$ ).

**Conclusion:** Our study revealed that essential oils from seeds and oleo-gum-resin of *Ferula assa-foetida* have antimicrobial properties. More laboratory studies are required to reach a definitive conclusion.

**Keywords:** *Ferula Foetida*; *Streptococcus Mutans*; *Streptococcus Sanguis*; *Streptococcus Salivarius*; *Streptococcus Sobrinus*; *Lactobacillus Rhamnosus*

- **Cite this article as:** Daneshkazemi A, Zandi H, Davari A, Vakili M, Emtiazi M, Lotfi R, Masoumi SMR. Antimicrobial Activity of the Essential Oil Obtained from the Seed and Oleo-Gum-Resin of *Ferula Assa-Foetida* against Oral Pathogens. *Front Dent*. 2019;16(2):113-120. doi: 10.18502/fid.v16i2.1362

## INTRODUCTION

Dental caries is the main reason for tooth loss and dental infections. This disease, which is

caused by several factors [1], begins with the accumulation of plaque on dental surfaces [2]. Poor oral health leads to an increase in the

accumulation of microorganisms and their derivatives. *Streptococcus mutans* (*S. mutans*) is the main bacterium found in dental plaque, which is colonized at the age of 6 to 9 months in the human's mouth [2,3] and causes damage to hard dental tissues through the fermentation of sucrose and the production of lactic acid [4]. This bacterium exploits sucrose to form dental plaque. The produced acid demineralizes hard dental tissues, while other bacteria in the *Streptococcaceae* family, such as *S. sobrinus*, *S. sanguis*, and *S. salivarius*, as well as *Lactobacilli*, can grow in an environment made by *S. mutans* and contribute to the development of tooth decay [5-7]. The nature of dental plaque is such that the plaque must be controlled regularly and effectively [8,9]. One of the methods for controlling the plaque is the use of mouthwashes [10]. Mouthwashes are particularly beneficial for patients with particular diseases, the elderly with diseases leading to weakening of the hand muscles, or people at risk of microbial infection, such as those susceptible to endocarditis or immune system weakness [11,12]. Mouthwashes can be taken daily and weekly [13]. Considering the increasing use of mouthwashes, it is essential to have access to substances with minimum side effects (pigment formation on dental surfaces, infections, and tissue toxicity) and maximum beneficial effects (plaque control) [14,15]. The scientific implementation of medicinal herbs has become possible with the advancement of chemistry and pharmacy. Nowadays, scientists prefer herbal remedies in many cases. *Ferula assa-foetida* is one of the plants belonging to the *Apiaceae* family [16,17]. This plant is under cultivation in different parts of the world, including the central and southern regions of Iran, and has been used in different countries [18]. Various studies have pointed to the antioxidant [19], antifungal [20,21], antiviral [22,23], anti-diabetic, and antihypertensive effects of this plant [24,25]. One of the important properties of this plant is its antimicrobial effect, which is a controversial issue. Fani et al [26] suggested that this plant is not effective against *S. mutans*, while studies by Rahman et al [27], Kavooosi et al [19], Kavooosi and Rowshan [28], and Haghghati et al [29] have reported the antimicrobial role of this plant against other bacteria such as *Staphylococcus aureus* and *Bacillus subtilis*. The discrepancies among previous studies

encouraged us to design a study aiming at investigating the antimicrobial effect of essential oils obtained from seeds and oleo-gum-resin of *Ferula assa-foetida* on oral pathogens including *S. mutans*, *S. sanguis*, *S. salivarius*, *S. sobrinus*, and *Lactobacillus rhamnosus* (*L. rhamnosus*).

## MATERIALS AND METHODS

### **Preparation of oleo-gum-resin essential oil:**

*Ferula assa-foetida* species, gathered from Tabas, Yazd Province, Iran during summer, was identified botanically by the Yazd Agricultural Research Center. The voucher number of the specimen was 2365. Then, 100 g of oleo-gum-resin was dissolved in one liter of distilled water to extract *assa-foetida* essential oil using hydrodistillation in the Clevenger device for 3 hours. After drying over anhydrous sodium sulfate, the attained essential oil was kept at 4°C until experimentation [30]. Finally, four different concentrations (2.5, 5, 10, and 20 µg/ml) of essential oil obtained from oleo-gum-resin were prepared.

### **Preparation of essential oil from *Ferula assa-foetida* seeds:**

First, 200 g of *Ferula assa-foetida* seeds were powdered and poured into 500 ml of double-distilled water. The hydrodistillation technique using the Clevenger device was employed to extract the seed essential oil which was protected from the light by being kept in glass containers hermetically sealed with rubber lids and covered with aluminum foil at 4°C.

### **Disk diffusion method for antimicrobial screening:**

The antimicrobial screening by seed and oleo-gum-resin essential oils of *Ferula assa-foetida* was conducted on standard strains of *S. mutans* (PTCC1683), *S. sanguis* (PTCC1449), *S. sobrinus* (PTCC1601), *S. salivarius* (PTCC1448), and *L. rhamnosus* (PTCC1637) which were prepared from the Iranian Research Organization for Science and Technology (IROST), Tehran, Iran. The antimicrobial screening was performed using the standard disk diffusion method in accordance with the Clinical and Laboratory Standards Institute (CLSI) 2017 protocols [30]. Fresh *Streptococcus* colonies were cultured on blood agar supplemented with 5% defibrinated sheep blood agar medium (Merck KGaA, Darmstadt, Germany), while De Man, Rogosa, and Sharpe (MRS) agar medium (Merck KGaA, Darmstadt, Germany) was used to culture *L.*

*rhamnosus* for inoculum. A turbidity equivalent to that of 0.5 McFarland turbidity standard, holding  $1.5 \times 10^8$  colony-forming units per milliliter (CFU/ml), was adjusted by inoculating fresh colonies into 5 ml of Mueller-Hinton broth (Merck KGaA, Darmstadt, Germany) followed by inoculation of *Streptococci* and *L. rhamnosus* using a sterile swab on Mueller-Hinton agar enriched with 5% defibrinated sheep blood agar medium (Merck KGaA, Darmstadt, Germany) and on MRS agar, respectively. Sterilized 6-mm blank paper disks (Padtan Teb Co., Tehran, Iran) were individually impregnated with 10  $\mu$ l of different concentrations (2.5, 5, 10, and 20  $\mu$ g/ml) of the essential oils, which were then cultured aseptically on a Mueller-Hinton agar and were incubated at 35°C with 5% carbon dioxide (CO<sub>2</sub>) for 48 hours. Afterward, the diameter (mm) of the non-growth zone was measured triplicate by a ruler to calculate the mean diameter. The controls consisted of disks impregnated with 0.2% Chlorhexidine and sterilized distilled water.

#### **Gas chromatography-mass spectrometry (GC-MS) analysis:**

The Hewlett-Packard 5971 GC-MS device (Avondale, PA, USA), available at the Islamic Azad University, Isfahan (Khorasgan) Branch, was used for analyzing the chemical composition of the essential oils with the following settings: 0.25 mm  $\times$  30 m polydimethylsiloxane DB-1 fused silica capillary column, 0.10- $\mu$ m film thickness, 1-ml/minute carrier gas of helium, injector temperature of 250°C, and detector temperature of 200°C. The column temperature was set variable from 35°C/minute to 180°C/minute at 4°C V/minute followed by 180°C/minute to 280°C/minute at 20°C V/minute. The electronic impact of 70 eV was considered for the mass spectra. A library computer search introduced the ingredients with their retention indices and visual interpretation of the mass spectra [31].

#### **Statistical Analysis:**

The data were analyzed using analysis of variance (ANOVA) and Kruskal-Wallis test in SPSS 17 software (SPSS Inc., Chicago, IL, USA).  $P < 0.05$  was considered as statistically significant.

## **RESULTS**

#### **Chemical compositions:**

The GC-MS findings exhibited that the main compounds found in the essential oil yielded from seeds of *Ferula assa-foetida* are alpha-D-

Xylofuranoside and methyl 2, 5-di-O-methyl- (30.2%), (E) -1-propenyl sec-butyl disulfide (13.13%), and (Z) -1-propenyl sec-butyl disulfide (11.34%; Table 1). In addition, the essential oil obtained from *Ferula assa-foetida* oleo-gum-resin mainly contained (E) -1-propenyl sec-butyl disulfide (36.15%) and (Z) -1-propenyl sec-butyl disulfide (27.93%; Table 2).

**Table 1:** Chemical composition of essential oil obtained from seeds of *Ferula assa-foetida*

No.	Composition	%
1	Alpha.-D-Xylofuranoside, methyl 2,5-di-O-methyl-	30.2
2	E-1-propenyl sec-butyl disulfide	13.13
3	Z-1-propenyl sec-butyl disulfide	11.34
4	Trifluoromethyl t-butyl disulfide	6.33
5	Disulfide, bis(1-methylpropyl)	5.47
6	10-epi-.gamma.-eudesmol	4.37
7	3-Mercapto propionitrile	3.28
8	Agarospinol	3.50
9	Ethanethioamide	2.90
10	Methyl sec-butyl disulphide	2.75
11	5-epi-7-epi- $\alpha$ -Eudesmol	2.62
12	1-(Methylthio) propyl propyl disulfide	2.53
13	(-)-Aristolene	1.41
14	(Z)-1-(But-2-en-1-yl)-2-(sec-butyl)disulfane	1.30
15	.alpha.-Pinene	1.80
16	3-thione-1,2-Dithiole	1.90
17	N-propyl sec-butyl disulfide	0.44
18	4-(hexadecyloxy)-3-nitrobenzenesulfonyl fluoride	0.39
19	Thiopivalic acid	0.36
20	Benzenepropanoic acid, pentyl ester	0.33
21	2-Thiazolidinethione	0.27
22	Morpholine, 2,6-dimethyl-	0.25
23	Elemol	0.17
24	Dimethyl trisulfide	0.17
<b>Total</b>		<b>94.03</b>

#### **Antibacterial activity:**

The antibacterial activity of essential oils yielded from seeds and oleo-gum-resin of *Ferula assa-foetida* plant was assessed by measuring the average growth inhibition zone. In the current study, the antimicrobial effects of four different concentrations (2.5, 5, 10, and 20  $\mu$ g/ml) of essential oils obtained from seeds and oleo-gum-resin were evaluated. The results indicated that both essential oils had significant antimicrobial activity at all concentrations ( $P < 0.05$ ). Both essential oils showed

**Table 2:** Chemical composition of essential oil obtained from oleo-gum-resin of *Ferula assa-foetida*

No.	Composition	%
1	E-1-propenyl <i>sec</i> -butyl disulfide	36.15
2	Z-1-propenyl <i>sec</i> -butyl disulfide	27.93
3	Guaiol	5.50
4	Carotol	5.14
5	bis (1-methyl propyl) disulfide	3.17
6	$\alpha$ -gurjunene	2.49
7	bis [(1-methylthio) propyl] disulfide	1.12
8	$\alpha$ - longipinene	1.86
9	Methyl penthyl tetrasulfide	1.16
10	Eudesmol(10-epi-gama)	0.98
11	Propyl <i>n</i> -butyl disulfide	0.97
12	$\delta$ - cadinene	0.95
13	Germacrene B	0.90
14	Methyl 1-(methylthio) ethyl disulfide	0.87
15	E- $\beta$ -ocimene	0.70
16	Methyl 1-(methylthio) propyl disulfide	0.66
17	$\beta$ - humulene	0.47
18	$\beta$ - himachalene	0.44
19	Valencene	0.36
20	Methyl <i>sec</i> -butyl disulfide	0.33
21	Eudesmol(7-epi-alpha)	0.33
22	Patchouli alcohol	0.29
23	$\alpha$ - humulene	0.28
24	Cedrene	0.24
25	$\alpha$ - ylangene	0.22
26	Longifolene	0.20
27	$\alpha$ - copaene	0.19
28	E- $\alpha$ - bisabolene	0.18
29	Cuparene	0.17
30	Neryl acetate	0.16
31	Spathulenol	0.15
32	allo-aromadendrene	0.13
32	$\beta$ -caryophyllene	0.12
34	Limonene	0.12
35	Z- $\beta$ -ocimene	0.12
36	p-cymene	0.10
37	Germacrene D	0.07
<b>Total</b>		<b>95.22</b>

antibacterial properties similar to that of Chlorhexidine (Tables 3 and 4).

The average growth inhibition zone was significantly dependent on the essential oil concentration for all bacteria ( $P < 0.05$ ) as the 20  $\mu\text{g/ml}$  concentration of both essential oils formed the largest average growth inhibition zone. Furthermore, this study showed that the essential oil from *Ferula assa-foetida* oleo-gum-resin had

significantly stronger antibacterial properties compared to the essential oil from *Ferula assa-foetida* seeds ( $P < 0.001$ ; Tables 3 and 4).

## DISCUSSION

Pathogenic microorganisms are one of the main health concerns for human beings. These microorganisms can be controlled easily by several antimicrobial chemicals; however, these chemicals can pose unwanted adverse complications [32]. Accordingly, there is a serious need for a safer therapeutic approach. Medicinal herbs are remarkable candidates and suitable alternatives to allopathic drugs. With the use of medicinal herbs, disadvantages of chemical drugs, such as high treatment costs, side effects, and development of resistance against antibiotics, can be avoided. Green medications have attracted a lot of attention, and the era of synthetic drugs is almost over. Effective ingredients derived from plants have been incorporated into official health care systems [32,33].

Iran is a country with various medicinal plants because of its unique and diverse geographical and climatic conditions; many of these herbs have shown positive therapeutic impacts such as antibacterial activity [34,35].

### *Chemical compositions:*

In the current study, the GC-MS results indicated that the main contents of seed essential oil are Alpha.-D-Xylofuranoside, methyl 2,5-di-O-methyl (30.2%), (E) -1-propenyl *sec*-butyl disulfide (13.13%), and (Z) -1-propenyl *sec*-butyl disulfide (11.34%). The contents of the essential oil obtained from oleo-gum-resin are (E) -1-propenyl *sec*-butyl disulfide (36.15%) and (Z) -1-propenyl *sec*-butyl disulfide (27.93%; Table 1).

These results are similar to those reported by Kavooosi et al [19], Khajeh et al [36], and Yousefi et al [37] although the levels of the compounds are slightly different. In the study by Kavooosi et al [19], the major components included (E) -1-propenyl *sec*-butyl disulfide (23.9%), 10-epi-c-eudesmol (15.1%), (Z) -1-propenyl *sec*-butyl disulfide (8.0%), (Z) - $\beta$ -ocimene (5.6%), and a-eudesmol (4.5%). In the study by Yousefi et al [37], the main compounds consisted of (E) -1-propenyl *sec*-butyl disulfide (50%), (Z) -1-propenyl *sec*-butyl disulfide (9.4%), globulol (12.5%), and a-pinene (8.8%). In the study by Khajeh et al [36], the major components were (E) -1-propenyl *sec*-butyl disulfide (40%), (Z) -



**Table 3:** Antibacterial effect of *Ferula assa-foetida* oleo-gum-resin essential oil on different bacteria based on inhibition zone diameter (mm), using the disk diffusion method

Species	Concentrations ( $\mu\text{g/ml}$ )*				Chlorhexidine 0.2%	P-value
	2.5	5	10	20		
<i>S. mutans</i>	13.5 $\pm$ 0.10	13.03 $\pm$ 0.58	14.07 $\pm$ 0.58	15.1 $\pm$ 0.10	15 $\pm$ 0.00	0.009
<i>S. sanguis</i>	7.07 $\pm$ 0.98	12.0 $\pm$ 0.00	12.17 $\pm$ 0.15	13.0 $\pm$ 0.00	15.53 $\pm$ 0.57	0.009
<i>S. sobrinus</i>	9.10 $\pm$ 0.10	11.2 $\pm$ 0.20	11.6 $\pm$ 0.62	12.13 $\pm$ 0.58	15.07 $\pm$ 0.12	0.011
<i>S. salivarius</i>	13.07 $\pm$ 0.57	12.08 $\pm$ 0.58	13.14 $\pm$ 0.15	14.03 $\pm$ 0.20	15.07 $\pm$ 0.57	0.009
<i>L. rhamnosus</i>	12.3 $\pm$ 0.10	13.16 $\pm$ 0.15	16.03 $\pm$ 0.57	18.1 $\pm$ 0.10	14.1 $\pm$ 0.10	0.009

\*The inhibition zone diameters (mm) were measured triplicate to report the mean values. Chlorhexidine was the positive control in this test as the reference antimicrobial agent.

**Table 4:** Antibacterial effect of essential oil obtained from seeds of *Ferula assa-foetida* on different bacteria based on inhibition zone diameter (mm), using the disk diffusion method

Species	Concentrations ( $\mu\text{g/ml}$ )*				Chlorhexidine 0.2%	P-value
	2.5	5	10	20		
<i>S. mutans</i>	8.87 $\pm$ 0.21	10.87 $\pm$ 0.21	12.07 $\pm$ 0.11	13.17 $\pm$ 0.15	15 $\pm$ 0.00	0.009
<i>S. sanguis</i>	6.94 $\pm$ 0.11	7.93 $\pm$ 0.30	9.90 $\pm$ 0.2	11.94 $\pm$ 0.20	15.53 $\pm$ 0.57	0.009
<i>S. sobrinus</i>	7.93 $\pm$ 0.15	10.16 $\pm$ 0.15	11.03 $\pm$ 0.2	10.06 $\pm$ 0.11	15.07 $\pm$ 0.12	0.013
<i>S. salivarius</i>	7.97 $\pm$ 0.38	9.0 $\pm$ 0.10	10.04 $\pm$ 0.058	12.0 $\pm$ 0.17	15.07 $\pm$ 0.57	0.012
<i>L. rhamnosus</i>	6.10 $\pm$ 0.10	7.13 $\pm$ 0.15	8.43 $\pm$ 0.20	8.8 $\pm$ 0.10	14.1 $\pm$ 0.10	0.011

\*The inhibition zone diameters (mm) were measured triplicate to report the mean values. Chlorhexidine was the positive control in this test as the reference antimicrobial agent

1-propenyl sec-butyl disulfide (8.7%), germacrene B (7.8%), and  $\alpha$ -pinene (5.9%). It should be noted that the current study was the first to evaluate the contents of *Ferula assa-foetida* seeds using the GC-MS, while other studies have only assessed the contents of oleo-gum-resin using the GC-MS.

This difference in the major compounds of *Ferula assa-foetida* essential oils can be related to the climatic conditions, the species used, the harvest time, and the method of processing.

#### **Antibacterial activity:**

In the current study, antimicrobial effects of four different concentrations (2.5, 5, 10, and 20  $\mu\text{g/ml}$ ) of essential oils obtained from seeds and oleo-gum-resin of *Ferula assa-foetida* on four oral bacteria were evaluated. The results indicated that all the concentrations had significant antimicrobial effects (Tables 3 and 4). Also, this study showed that this effect can be similar to that of Chlorhexidine. These results are similar to the results reported by Kavooosi and Rowshan [28] and Siddiqui et al [38] although these two studies have evaluated the antimicrobial effect of essential oil obtained from *Ferula* oleo-gum-resin on *Staphylococcus aureus*, *Bacillus Subtilis*, and *Escherichia Coli*.

Both studies used the minimum inhibitory concentration (MIC) method, whereas the disk diffusion method was used in the current study.

The results of the current study are not similar to the results of the study by Fani et al [26], which showed that *Ferula assa-foetida* does not have any antimicrobial effect on *S. mutans* and *S. sanguis*.

The cited study evaluated aqueous and ethanolic extracts of *Ferula*, while the current study evaluated essential oils obtained from *Ferula assa-foetida* seeds and oleo-gum-resin. The mentioned study used the well diffusion method and MIC, while the current study implemented the disk diffusion method. The cited study did not consider any of the concentrations effective on these two bacteria, while in the current study, the lowest assessed concentration (2.5  $\mu\text{g/ml}$ ) resulted in a growth inhibition zone with a diameter of 8.87 $\pm$ 0.21 mm with seed essential oil and a diameter of 13.5 $\pm$ 0.1 mm with oleo-gum-resin essential oil against *S. mutans*. It seems that the difference in the results is caused by different extracts of *Ferula assa-foetida* and the method used for measuring the antimicrobial effect. Kavooosi and Rowshan [28] investigated the antioxidant and antibacterial activities of *Ferula assa-foetida*

and showed that this plant has inhibitory effects on the growth of gram-positive bacteria ( $P < 0.05$ ). Siddiqui et al [38] reported dose-dependent antimicrobial effects of *Ferula assa-foetida* essential oil on gram-positive bacteria; the 50 µg/ml concentration had no significant antibacterial effect compared to standard antibiotics, whereas a concentration of 100 µg/ml was more effective compared to standard antibiotics. This result is in line with the findings of the study by Haghghati et al [29] and our study. Based on the reports by Fani et al [26], there was no significant difference in the antibacterial effects of *Ferula assa-foetida* and *Quercus infectoria* extracts when comparing *S. mutans* and *S. sanguis*. According to the antimicrobial properties of essential oils obtained from medicinal plants, phenolic monoterpenes are the most important effective substances that can inhibit bacterial growth due to increased permeability and polarizability of the cell membrane, resulting in microleakage of protons from the cells, electric potential imbalance in the membrane, proton motive force reduction, and decreased adenosine triphosphate (ATP) formation [39]. It should be noted that the consequence of decreased membrane potential is the loss of ions, ATP, amino acids, and proteins following the leakage from the cell [40]. One of the indications of membrane damage and cell death is the ion leakage out of the cell [40]. Extracts of several *Ferula* species have been shown to have moderate antibacterial activity, probably due to the presence of phenols, flavonoids, and sesquiterpenes. High levels of phenolic and flavonoid compounds are associated with the antibacterial activity [41,42].

The hydrophobicity and solubility of the lipid in the compounds can be the reason for their antibacterial activity. However, there are various mechanisms for the antibacterial features of different essential oils. Extensive diffusion of compounds into the lipid bilayer and increased membrane permeability induced by some substances have revealed potent antibacterial activity. Acyclic sulfur-containing compounds are mainly found in both essential oils [43]. Kavooosi et al [19] reported a more effective antimicrobial activity for cyclic compounds compared to acyclic sulfur-containing compounds. The diffusion into the lipid bilayer and membrane permeability are significantly elevated by cyclic compounds

through their higher spatial volume, resulting in high membrane permeability and cell death [43].

Nevertheless, more comprehensive studies are required to draw a definitive conclusion regarding the antibacterial activity of medicinal plants and their effective concentrations. Active collaboration of research institutes in different parts of the world is needed to accelerate the investigations in this regard.

## CONCLUSION

The present study revealed that essential oils obtained from seeds and oleo-gum-resin of *Ferula assa-foetida* show antimicrobial effects against oral pathogens.

## CONFLICT OF INTEREST STATEMENT

None declared.

## REFERENCES

1. Murray PE, About I, Franquin JC, Remusat M, Smith AJ. Restorative pulpal and repair responses. *J Am Dent Assoc.* 2001 Apr;132(4):482-91.
2. Karikalan S, Mohankumar A. Studies on ampicillin resistant plasmid of *Streptococcus mutans* isolated from dental caries patients. *Biosci Biotech Res Comm.* 2016;9(1):151-6.
3. Loesche WJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev.* 1986 Dec;50(4):353-380.
4. Yadav K, Prakash S. Dental caries: a review. *Asian J Biomed Pharm Sci.* 2016;6(53):1-7.
5. Brännström M, Nordenvall KJ. Bacterial penetration, pulpal reaction and the inner surface of concise enamel bond. Composite fillings in etched and unetched cavities. *J Dent Res.* 1978 Jan;57(1):3-10.
6. Cox CF, Keall CL, Keall HJ, Ostro E, Bergenholtz G. Biocompatibility of surface-sealed dental materials against exposed pulps. *J Prosthet Dent.* 1987 Jan;57(1):1-8.
7. Westergren G, Emilson CG. Colonization and cariogenic potential in hamsters of the bacterium *Streptococcus sanguis* isolated from human dental plaque. *Arch Oral Biol.* 1982;27(10):817-22.
8. Bjørndal L, Darvann T, Thylstrup A. A quantitative light microscopic study of the odontoblast and subodontoblastic reactions to active and arrested enamel caries without cavitation. *Caries Res.* 1998 Jan-Feb;32(1):59-69.
9. Brännström M, Lind PO. Pulpal response

- to early dental caries. *J Dent Res.* 1965 Sep-Oct;44(5):1045-50.
10. Browning WD, Johnson WW, Gregory PN. Clinical performance of bonded amalgam restorations at 42 months. *J Am Dent Assoc.* 2000 May;131(5):607-11.
  11. Francis JR, Hunter B, Addy M. A comparison of three delivery methods of chlorhexidine in handicapped children: I. Effects on plaque, gingivitis, and toothstaining. *J Periodontol.* 1987 Jul;58(7):451-5.
  12. Kalaga A, Addy M, Hunter B. Comparison of chlorhexidine delivery by mouthwash and spray on plaque accumulation. *J Periodontol.* 1989 Mar;60(3):127-30.
  13. Atkinson CB. Hints, queries and comments: pyrozone. *Dent Cosmos.* 1893;35:330-2.
  14. Patel R, Gallagher J, Chapple I. Question from practice: how to select the right mouthwash. *J Cereb Circ.* 2018;13:57.
  15. Amirzade-Iranaq MH, Masoumil SMR. 50: effect of low-level laser therapy in the treatment of burning mouth syndrome: a systematic review and meta-analysis. *BMJ Open.* 2017;7(Suppl 1):bmjopen-2016-015415.50.
  16. Saeidy S, Nasirpour A, Keramat J, Desbrières J, Le Cerf D, Pierre G, et al. Structural characterization and thermal behavior of a gum extracted from *Ferula assa foetida* L. *Carbohydr Polym.* 2018 Feb 1;181:426-32.
  17. Kasaian J, Mohammadi A. Biological activities of farnesiferol C: a review. *J Asian Nat Prod Res.* 2018 Jan;20(1):27-35.
  18. Zomorodian K, Saharkhiz J, Pakshir K, Immeripour Z, Sadatsharifi A. The composition, antibiofilm and antimicrobial activities of essential oil of *Ferula assa-foetida* oleo-gum-resin. *Biocatal Agric Biotechnol.* 2018 Apr;14:300-304.
  19. Kavooosi G, Tafsiry A, Ebdam AA, Rowshan V. Evaluation of antioxidant and antimicrobial activities of essential oils from *Carum copticum* seed and *Ferula assafoetida* latex. *J Food Sci.* 2013 Feb;78(2):T356-61.
  20. Angelini P, Pagiotti R, Venanzoni R, Granetti B. Antifungal and allelopathic effects of *Asafoetida* against *Trichoderma harzianum* and *Pleurotus* spp. *Allelopathy J.* 2009 Apr;23(2):357-68.
  21. Gowda NKS, Malathi V, Suganthi RU. Effect of some chemical and herbal compounds on growth of *Aspergillus parasiticus* and aflatoxin production. *Anim Feed Sci Technol.* 2004 Oct;116(3-4):281-91.
  22. Lee CL, Chiang LC, Cheng LH, Liaw CC, Abd El-Razek MH, Chang FR, et al. Influenza A (H1N1) antiviral and cytotoxic agents from *Ferula assa-foetida*. *J Nat Prod.* 2009 Sep;72(9):1568-72.
  23. Rollinger JM, Steindl TM, Schuster D, Kirchmair J, Anrain K, Ellmerer EP, et al. Structure-based virtual screening for the discovery of natural inhibitors for human rhinovirus coat protein. *J Med Chem.* 2008 Feb 28;51(4):842-51.
  24. Esmaeili H, Hafezimoghadam Z, Esmailidehaj M, Rezvani ME, Hafizibarjin Z. The effect of asafoetida essential oil on myocardial ischemic-reperfusion injury in isolated rat hearts. *Avicenna J Phytomed.* 2018 Jul-Aug;8(4):338-349.
  25. Al-Ja'fari AH, Vila R, Freixa B, Costa J, Cañigüeral S. Antifungal compounds from the rhizome and roots of *Ferula hermonis*. *Phytother Res.* 2013 Jun;27(6):911-5.
  26. Fani MM, Bazargani A, Farboodnia MA, Hasanpour Z, Zamani K, Yousefi Manesh E. An in Vitro Study on the Antibacterial Effect of *Ferula Assa-Foetida* L. and *Quercus Infectoria* Olivier Extracts on *Streptococcus Mutans* and *Streptococcus Sanguis*. *Avicenna J Dent Res.* 2015 Jun;7(1):10.17795/ajdr-22656.
  27. Rahman MU, Gul S, Odhano EA. Antimicrobial activities of *Ferula assafoetida* oil against gram positive and gram negative bacteria. *Am-Euras J Agric Environ Sci.* 2008 Jan;4(2):203-6.
  28. Kavooosi G, Rowshan V. Chemical composition, antioxidant and antimicrobial activities of essential oil obtained from *Ferula assa-foetida* oleo-gum-resin: effect of collection time. *Food Chem.* 2013 Jun 15;138(4):2180-7.
  29. Haghghati F, Jafari S, Beyt Elahi JM. Comparison of antimicrobial effects of ten herbal extracts with Chlorhexidine on three different oral pathogens; an in vitro study. *Hakim Res J.* 2003 Fall;6(3):71-6.
  30. Bagheri SM, Asl AA, Shams A, Mirghanizadeh-Bafghi SA, Hafizibarjin Z. Evaluation of cytotoxicity effects of oleo-gum-resin and its essential oil of *Ferula assa-foetida* and ferulic acid on 4T1 breast cancer cells. *Indian J Med Paediatr Oncol.* 2017 Apr-Jun;38(2):116-20.
  31. Botelho MA, Nogueira NA, Bastos GM, Fonseca SG, Lemos TL, Matos FJ, et al. Antimicrobial activity of the essential oil from *Lippia sidoides*, carvacrol and thymol against oral pathogens. *Braz J Med Biol Res.* 2007 Mar;40(3):349-56.

32. Ahmad I, Beg AZ. Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *J Ethnopharmacol.* 2001 Feb;74(2):113-23.
33. Khan R, Islam B, Akram M, Shakil S, Ahmad A, Ali SM, et al. Antimicrobial activity of five herbal extracts against multi drug resistant (MDR) strains of bacteria and fungus of clinical origin. *Molecules.* 2009 Feb 4;14(2):586-97.
34. Dip EC, Pereira NA, Fernandes PD. Ability of eugenol to reduce tongue edema induced by *Dieffenbachia picta* Schott in mice. *Toxicol.* 2004 May;43(6):729-35.
35. Dehpour AA, Ebrahimzadeh MA, Nabavi SF, Nabavi SM. Antioxidant activity of the methanol extract of *Ferula assafoetida* and its essential oil composition. *Grasas y Aceites.* 2009 Sep;60(4):405-12.
36. Khajeh M, Yamini Y, Bahramifar N, Sefidkon F, Pirmoradei MR. Comparison of essential oils compositions of *Ferula assafoetida* obtained by supercritical carbon dioxide extraction and hydrodistillation methods. *Food Chem.* 2005 Aug;91(4):639-44.
37. Yousefi M, Mohammadi M, Habibi Z. Disulphides in the volatile oil of *Ferula behboudiana* Rech. f. & Esfand. *Nat Prod Res.* 2011 Oct;25(17):1629-34.
38. Siddiqui RR, Zafar U, Chaudhry SS, Ahmad H. Antimicrobial activity of essential oils from *Schinus terebinthifolius*, *Cypress sempervirens*, *Citrus limon*, *Ferula assafoetida*. Part I. *Pak J Sci Ind Res.* 1995;38(9/10):358-61.
39. García-García R, López-Malo A, Palou E. Bactericidal action of binary and ternary mixtures of carvacrol, thymol, and eugenol against *Listeria innocua*. *J Food Sci.* 2011 Mar;76(2):M95-100.
40. Rúa J, Fernández-Álvarez L, de Castro C, del Valle P, de Arriaga D, García-Armesto MR. Antibacterial activity against foodborne *Staphylococcus aureus* and antioxidant capacity of various pure phenolic compounds. *Foodborne Pathog Dis.* 2011 Jan;8(1):149-57.
41. Eftekhari F, Yousefzadi M, Borhani K. Antibacterial activity of the essential oil from *Ferula gummosa* seed. *Fitoterapia.* 2004 Dec;75(7-8):758-9.
42. Ibraheim ZZ, Abdel-Mageed WM, Dai H, Guo H, Zhang L, Jaspars M. Antimicrobial antioxidant daucane sesquiterpenes from *Ferula hermonis* Boiss. *Phytother Res.* 2012 Apr;26(4):579-86.
43. Dorman H, Deans S. Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *J Appl Microbiol.* 2000 Feb;88(2):308-16.