

Labial Ankyloglossia Associated with Oligodontia: A Case Report

L. Chandrashekar¹, K. R. Kashinath², Setty Suhas³

¹Reader, Department of Oral Medicine and Radiology, Best Dental Science College and hospital Madurai, Tamilnadu, India

²Principal, Professor and Head, Department of Prosthodontics, Sri Siddhartha Dental College and Hospital, Tumkur, India

³Professor, Department of Oral Medicine and Radiology, Sri Siddhartha Dental College and Hospital, Tumkur, India

✉ Corresponding author:

L. Chandrashekar, Department of Oral Medicine and Radiology, Best Dental Science College and hospital Madurai, Tamilnadu, India

drshekar75@gmail.com

Received: 13 February 2014

Accepted: 17 April 2014

Abstract

Developmental disturbances of the tongue (aglossia, macroglossia, microglossia, and ankyloglossia) can adversely affect the development of the surrounding structures including the palate, alveolar process, and teeth. These developmental disturbances impair functions such as mastication, speech, and swallowing. Ankyloglossia is of various types and may be associated with other syndromic features like cleft palate. We report a rare type of non-syndromic ankyloglossia associated with missing of the permanent anterior tooth.

Key Words: Ankyloglossia; Oligodontia; Labial Frenum

Journal of Dentistry, Tehran University of Medical Sciences, Tehran, Iran (2014; Vol. 11, No. 4)

INTRODUCTION

Ankyloglossia is the restriction of tongue movement due to congenital abnormality in the lingual frenum. Ankyloglossia may be syndrome associated, but commonly it has a non-syndromic presentation with an incidence of up to 10% [1].

We present a case with an unusual type of ankyloglossia that cannot be classified using the current classification system for ankyloglossia, wherein only the tip of the tongue is fused with the labial frenum of the mandibular arch and is associated with missing of the mandibular permanent anterior teeth.

CASE REPORT

A 13-year-old girl was seen for complaining of decayed teeth. This was her first dental visit.

On examination, the mandibular right lateral incisors and bilateral central incisors were missing and the left mandibular lateral incisor was decayed. She also had generalized enamel hypoplasia and angle's class I malocclusion with bimaxillary protrusion. Her tongue was depapillated and ankylosis of the tip of the tongue with the labial frenum of the lower lip was noticed (Figure 1).

Movements of the tongue were restricted and pronunciations of consonants were affected. Family history was unremarkable. A pediatrician evaluated her general health status and she was found to be suffering from iron deficiency anemia.

After taking detailed history and complete examination, a provisional diagnosis of labial ankyloglossia associated with oligodontia and iron deficiency anemia was considered.



Fig 1. Fusion of tip of tongue with labial frenum

Panoramic radiography confirmed missing of the permanent mandibular central incisors and the right lateral incisor (figure 2).

After routine blood investigations, surgical release of the ankyloglossia with “Z” plasty using a 15 no. B.P blade was carried out under local anesthesia and was sutured using vicryl 3-0 suture material. Post treatment healing was uneventful and tongue movement was restored to normalcy (figure 3). Iron deficiency anemia was managed by the pediatrician uneventfully.

DISCUSSION

Treatment seeking behavior for structural birth defects with minimal or no functional deficit are generally delayed by patients, more so in the lower socio economic strata of the society. In the present case, parents of the patient had not sought any consultation for their child’s tongue problem because they felt that no treatment was essential as her daily activities were not affected. Our patient presented for treatment of a decayed tooth and the presence of ankyloglossia and oligodontia were identified.

Ankyloglossia is defined as “a condition in which the tip of the tongue cannot be protruded beyond the lower incisor teeth because of short frenulum linguae, often containing scar tissue” [2]. Lower lip develops by the




Fig 2. OPG showing missing mandibular central incisors and left lateral incisor

fusion of bilateral mandibular arches at Carnegie stage 11 of embryogenesis and it completely merges by stage 15 and the tongue tissue is found to be prominent by stage 16 [3]. In the present case, fusion of the tip of the tongue to the lower lip and missing of the permanent teeth suggest that there has been a deviation from the normal development in the region of fusion of the first brachial arches. On the review of literature, various classifications of ankyloglossia have been proposed [4]; we were unable to classify our case based on the published criteria. Incidental findings associated with ankyloglossia including median mandibular clefts [5], bifid uvula [6], undeveloped epiglottis or hyoid and underdeveloped thyroid cartilage [7] have been reported. Ankyloglossia has also been observed together with syndromes such as X-linked cleft palate syndrome [8] Kindler syndrome [9], Vander Woude syndrome [10] and Opitz syndrome [9]. Labial frenum ankyloglossia has also been reported in a child associated with oligodontia of the deciduous molar dentition [10]. In the present case, ankyloglossia was associated with partial anodontia of the permanent anterior teeth and this association has not been reported previously. Presence of iron deficiency anemia in the present case could be another non related finding. Hanhart syndrome and holoprosencephaly or microforms of holoprosencephaly have been associated with tongue anomalies; however, other features like skeletal abnormalities of the midface, limbs and CNS anomalies are also part of the spectrum



Fig 3 .Tongue after surgical release

of other anomalies in these patients [11, 12]. Abnormal development of the palate (cleft palate) and tongue (ankyloglossia) have been associated with aberrations in T-bx22 genes. Mutations in this gene have been associated with cleft palate and ankyloglossia, and it is believed to play a major role in human palatogenesis [13]. Preferential tooth agenesis (incisors and/or premolars) associated with orofacial clefts have been identified in Interferon Regulatory Factor 6 (IRF6) gene deletions and point mutations [14]. As there was no cleft palate in our case, we can draw an inference of the possibility of absence of this mutation.

REFERENCES

1- Segal LM, Stephenson R, Dawes M, Feldman P. Prevalence, diagnosis, and treatment of ankyloglossia: methodologic review. *Can Fam Physician*. 2007 Jun;53(6):1027-33.
 2- Wallace AF. Tongue tie. *Lancet*. 1963 Aug 24;2(7304):377-8.

3- Yoon H, In Chung IS, Seol EY, Park BY, Park HW. Development of the lip and palate in staged human embryos and early fetuses. *Yonsei Med J*. 2000 Aug;41(4):477-84.

4- Suter VG, Bornstein MM. Ankyloglossia: Facts and myths in diagnosis and treatment. Review. *J Periodontol*. 2009 Aug;80(8):1204-19.

5- McCarthy JG. Plastic surgery: cleft lip and palate and cranio-facial anomalies, Vol. 4. 3rd ed. Philadelphia: WB Saunders; 1990. p. 2922-41.

6- Forbes SA, Brennan L, Richardson M, Coffey A, Cole CG, Gregory SG, et al. Refined mapping and YAC contig construction of the X-linked cleft palate and ankyloglossia locus (CPX) includes the proximal X-V homology breakpoint within Xq21.3. *Genomics*. 1996 Jan 1;31(1):36-43.

7- Hacham-Zadeh S, Garfunkel AA. Kindler syndrome in two related Kurdish families. *Am J Med Genet*. 1985 Jan;20(1):43-8.

8- Burdick AB, Ma LA, Dai ZH, Gao NN. Van der Woude syndrome in two families in China. *J Craniofac Genet Dev Biol*. 1987;7(4):413-8.

9- Brooks JK, Leonard CO, Cocco PJ Jr. Opitz (BBB/ G) syndrome: Oral manifestations. *Am J Med Genet*. 1992 Jun 1;43(3):595-601.

10- Rana RE, Puri VA, Thakkur RK, Baliarsing AS. Median cleft of mandible and lower lip with ankyloglossia and ectopic minor salivary gland on tongue. *Indian J Plast Surg*. 2004;37(1):67-70.

11- Tüysüz B, Erginel A, Unutmaz T, Cenani A. Hypoglossia-hypodactylia (Hanhart's) syndrome with sensorineural hearing loss. *Turk J Pediatr*. 1994 Oct-Dec;36(4):347-52.

12- Kauvar EF, Solomon BD, Curry CJ, van Essen AJ, Janssen N, Dutra A, et al. Holoprosencephaly and agnathia spectrum: Presentation of two new patients and review of the literature.

rature. *Am J Med Genet C Semin Med Genet.* 2010 Feb 15;154C(1):158-69.

13- Andreou AM, Pauws E, Jones MC, Singh MK, Bussen M, Doudney K, et al. TBX22 missense mutations found in patients with X-linked cleft palate affect DNA binding, sumoylation, and transcriptional repression. *Am J Hum Genet.* 2007 Oct;81(4):700-12. Epub

2007 Aug 16.

14- Vieira AR, Modesto A, Meira R, Barbosa AR, Lidral AC, Murray JC. Interferon regulatory factor 6 (*IRF6*) and fibroblast growth factor receptor 1 (*FGFR1*) contribute to human tooth agenesis. *Am J Med Genet A.* 2007 Mar 15;143(6):538-45.